

Lecture 7: Stochastic models for small groups such as households


A Brief History of the Reed-Frost Model

- PD En'ko (1889) Deterministic difference equations
- L Reed and WH Frost (1930) Marbles and shoots
- M Greenwood (1931) Alternative formulation
- H Abbey (1952) 1st analysis as a stochastic process
- L Elveback, JP Fox, E Ackerman (1960) 1st computer program and lots of theory

Reed-Frost Model



Lowell Reed
1886 - 1966



Wade Hampton Frost
1880–1938

Both Former Deans: Johns Hopkins School of Public Health



Helen Abbey
1915 - 2001



Eugene Ackerman
1920 -



John P. Fox
Died around 1989

Reed-Frost Model

Stochastic process: discrete state space and time $t_0, t_1, t_2 \dots$

- Infectious agent natural history
 - ◆ Infectious for one time unit
- Social contact structure
 - ◆ Random mixing
 - ◆ $p = 1 - q$, probability two people make contact sufficient to transmit
- $R_0 = (n-1)p$

Reed-Frost Model

$$P(I_{t+1} | S_t, I_t) = \binom{S_t}{I_{t+1}} (1 - q^{I_t})^{I_{t+1}} q^{I_t(S_t - I_{t+1})}, S_t \geq I_{t+1},$$

$$S_{t+1} = S_t - I_{t+1},$$

$$R_{t+1} = R_t + I_t,$$

$$S_t + I_t + R_t = n, \forall t,$$

$$P[S(0) = n - 1] = 1, P[I(0) = 1] = 1, P[R(0) = 0] = 1$$

$\{S_t, I_t\}_{t=0,1,\dots}$ is a Markov chain

Greenwood Model

$$P(I_{t+1} | S_t, I_t) = \binom{S_t}{I_{t+1}} (1 - q^{I_t})^{I_{t+1}} q^{I_t (S_t - I_{t+1})}, S_t \geq I_{t+1},$$

$$S_{t+1} = S_t - I_{t+1},$$

$$R_{t+1} = R_t + I_t,$$

$$S_t + I_t + R_t = n, \forall t,$$

$$P[S(0) = n - 1] = 1, P[I(0) = 1] = 1, P[R(0) = 0] = 1$$

$\{S_t, I_t\}_{t=0,1,\dots}$ is a Markov chain

A Chain

$$I_0 \rightarrow I_1 \rightarrow I_2 \rightarrow \cdots \rightarrow I_r$$

$$\begin{aligned} P(I_0, I_1, I_2, \cdots, I_r) &= \\ &P(I_1 \mid S_0, I_0)P(I_2 \mid S_1, I_1) \cdots \\ &P(I_r \mid S_{r-1}, I_{r-1}) \end{aligned}$$


$$T = \inf_{t \geq 0} \{t : S_t I_t = 0\}.$$

Example

Possible individual chains when $S_0 = 3$, $I_0 = 1$


Chain	Probability	Final Size
$\{i_0, i_1, i_2, \dots, i_T\}$		R_T
$\{1\}$	q^3	1
$\{1, 1\}$	$3pq^4$	2
$\{1, 1, 1\}$	$6p^2q^4$	3
$\{1, 2\}$	$3p^2q^3$	3
$\{1, 1, 1, 1\}$	$6p^3q^3$	4
$\{1, 1, 2\}$	$3p^3q^2$	4
$\{1, 2, 1\}$	$3p^3q(1 + q)$	4
$\{1, 3\}$	p^3	4

Epidemic in the Community




Community Structure

Community




n - Households (Semiclosed Groups)

Definition of Time Intervals



Onset of Cases in Household




"IN ACCEPTING THE FAMILY AS A SEMICLOSED GROUP, ONE MUST BELIEVE THAT INFECTIONS IN SUCCESSIVE FAMILY MEMBERS WITHIN A SHORT PERIOD OF TIME ARE MORE LIKELY TO BE RELATED TO EACH OTHER THAN TO HAVE BEEN SEPERATELY ACQUIRED FROM OUTSIDE SOURCES. THIS IS ESSENTUALLY AN ACT OF FAITH AND NEARLY IMPOSSIBLE TO ESTABLISH IN TERMS OF EXACT PROBABILITY."

BUCK, C. ACUTE UPPER RESPIRATORY INFECTIONS IN FAMILIES
Am. J. Hyg. 63, 1-12, 1956.

1. NATURAL HISTORY OF S-I-R INFECTION PROCESS

USCEPT.
INFECT.
REMOVED



FLU: $E(T_1) \cong E(T_3) \cong 2$ DAYS
 $E(T_2) \cong 4$ DAYS

MEASLES: $E(T_1) \cong 8$ DAYS
 $E(T_3) \cong 9$ DAYS
 $E(T_2) \cong 4.5$

PROBABILITY MODEL FOR ACUTE INFECTIOUS DISEASES

Π_{1j} PROB. THAT 1 OUT OF j PERSONS ARE INFECTED, $1 \leq j$

ASSUMPTIONS:

1. T AND T_2 ARE FIXED ($B = b^t$ and $Q = q^{t_2}$)
2. RANDOM MIXING WITHIN HOUSEHOLDS
3. PROB. FAMILY MEMBER INFECTED FROM COMM.
IS INDEPENDENT OF # INFECTED IN HOUSEHOLD

3. TRANSMISSION PROBABILITIES

WITHIN FAMILIES:

p = PROB. INFECTED GIVEN CONTACT PER UNIT OF TIME

$$q = 1 - p$$

$$Q = E(q^{T^2}).$$

SAR = $(1 - Q) \times 100$ SECONDARY ATTACK RATE

FROM COMMUNITY SOURCES:

T = DURATION OF THE EPIDEMIC IN THE COMMUNITY

a = PROB. INFECTED FROM COMMUNITY SOURCES OVER

THE COURSE OF THE EPIDEMIC, $b = 1 - a$

$$B = E(b^T)$$

CPI = $1 - B$ COMMUNITY PROBABILITY OF INFECTION

PARTITION SAMPLE SPACE INTO i INFECTIVES AND $j - i$ ESCAPES.
WHERE THERE ARE $\binom{j}{i}$ SUCH PARTITIONS

FOR ONE PARTITION OF THE SAMPLE SPACE:

E - EVENT i OUT OF i INFECTED

$$P(E) = \pi_{ii}$$

F - EVENT $j - i$ ESCAPE

$$P(F|E) = B^{j-i} Q^{i(j-i)}, \quad i < j$$

THEN WE HAVE

$$\pi_{ij} = P(E \cap F) = P(E) P(F|E)$$

$$\pi_{ij} = \pi_{ii} B^{j-i} Q^{i(j-i)}, \quad i < j.$$

$$\pi_{jj} = 1 - \sum_{i=0}^{j-1} \pi_{ij}$$

SPECIAL CASES

BINOMIAL: WHEN $Q = 1$, THEN

$$\Pi_{ij} = \binom{j}{i} (1 - B)^i B^{j-i}, \quad i \leq j$$

REED-FROST-EN'KO MODEL: WHEN $B = 1$, AND THERE ARE $i_0 \geq 1$ INITIAL INFECTIVES, THEN

$$\Pi_{ij} = \binom{j}{i} \Pi_{i1} Q^{(i_0+1)(j-1)}, \quad i < j.$$

$$\Pi_{jj} = 1 - \sum_{i=0}^{j-1} \Pi_{ij}$$

BAILEY (1975). LUDWIG (1975)

"DISCORDANT COUPLES MODEL": WHEN $j = 1$, $i_0 = 1$, AND $B < 1$.

THEN $\Pi_{01} = BQ$ AND $\Pi_{11} = 1 - BQ$.

$$\pi_{ij} = \binom{j}{i} \pi_{11} B^{j-1} Q^{i(j-1)}, \quad i < j.$$




$$\pi_{jj} = 1 - \sum_{i=0}^{j-1} \pi_{ij}$$

LONGINI & KOOPMAN (1982). LONGINI, ET AL. (1982)

$$\pi_{01} = B$$

$$\pi_{11} = 1 - B$$

PROBABILITY MASS FUNCTION FOR A HOUSEHOLD WITH FIVE SUSCEPTIBLES



DISTRIBUTION OF RESPIRATORY SYMPTOMS
DURING H3N2 EPIDEMIC PERIOD OF 1977-78 IN TECUMSEH

NUMBER INFECTED	NUMBER OF INDIVIDUALS/HOUSEHOLD				
	1	2	3	4	5
0	38	48 (48)	13 (13)	6 (6)	4 (4)
1	19	29 (44)	19 (27)	13 (20)	3 (11)
2		25 (10)	20 (18)	17 (24)	9 (12)
3			10 (4)	23 (13)	8 (6)
4				7 (3)	5 (2)
5					6 (0)
TOTAL	57	102	62	66	35
	n_1	n_2	n_3	n_4	

2

LIKELIHOOD FUNCTION

n - Total number of households

n_j - Number of households with j initial
susceptibles

n_{ij} - Number of households with i of j
individuals infected

$$\left(\sum_i n_{ij} = n_j, \sum_j \sum_i n_{ij} = \sum_j n_j = n \right)$$

$$L(Q, B) = \prod_j \prod_i \pi_{ij}^{n_{ij}}$$

LONGINI & KOOPMAN (1982)

$$\hat{B} = \frac{1}{n} \sum_j n_{\cdot j} \left(\frac{n_{0j}}{n_{\cdot j}} \right)^{1/j} \sim 90\% \text{ EFF.}$$

DISTRIBUTION OF INFECTIONS BY HOUSEHOLDS

Number of Susceptibles/Household

#Infected	j				
1	2	3		J
0	n_{01}	n_{02}	n_{03} n_{0J}	
1	n_{11}	n_{12}	n_{13} n_{1J}	
2	-	n_{22}	n_{23} n_{2J}	
.	
.	
.	
.	
.	
J	-	-	- n_{JJ}	
	$n_{.1}$	$n_{.2}$	$n_{.3}$ $n_{.J}$	
					n

2:3

DISTRIBUTION OF INFLUENZA A(H3N2) INFECTIONS

1980-81 TECUMSEH DATA

Number of Susceptibles/Household

Number Infected	Number of Susceptibles/Household				Total
	1	2	3	4	
0	44	62	47	38	
1	10	13	8	11	
2	-	9	2	7	
3	-	-	3	5	
4	-	-	-	1	
	54	84	60	62	260
	n ₁	n ₂	n ₃	n ₄	

MAXIMUM LIKELIHOOD ESTIMATION

$$\ln L = c + \sum_j \sum_i n_{ij} \left\{ \ln \pi_{ij} + (j-i) \ln B + j(j-i) \ln Q \right\}$$

$\{0 \leq B \leq 1, 0 \leq Q \leq 1\}$

MLE'S \hat{Q} AND \hat{B} ARE SOLUTIONS OF

$$0 = \frac{\partial \ln L}{\partial Q} \Big|_{\hat{Q}, \hat{B}} = \sum_j \sum_i n_{ij} \left\{ \frac{1}{\pi_{ij}} \left(\frac{\partial \pi_{ij}}{\partial Q} \right) + \frac{j(j-i)}{Q} \right\}$$

$$0 = \frac{\partial \ln L}{\partial B} \Big|_{\hat{Q}, \hat{B}} = \sum_j \sum_i n_{ij} \left\{ \frac{1}{\pi_{ij}} \left(\frac{\partial \pi_{ij}}{\partial B} \right) + \frac{j-i}{B} \right\}$$

SOLVE BY METHOD OF SCORING WHICH

ALSO PROVIDES

$\hat{Var}(\hat{Q})$, $\hat{Var}(\hat{B})$ AND $\hat{Cov}(\hat{Q}, \hat{B})$.

ALSO $\hat{Q} \sim N$ AND $\hat{B} \sim N$ LARGE N .

LONGINI & KOOPMAN (1982)

INFORMATION MATRIX

$$-E\left(\frac{\partial^2 \ln L}{\partial Q^2}\right) = \sum_k \sum_j n_k m_{jk} \left\{ \frac{1}{m_{jj}} \left[\frac{1}{m_{jj}} \left(\frac{\partial m_{jj}}{\partial Q}\right)^2 - \frac{\partial^2 m_{jj}}{\partial Q^2} \right] + \left[\frac{j(k-j)}{Q^2} \right] \right\}$$

$$-E\left(\frac{\partial^2 \ln L}{\partial Q \partial B}\right) = \sum_k \sum_j n_k \left(\frac{m_{jk}}{m_{jj}}\right) \left[\frac{1}{m_{jj}} \left(\frac{\partial m_{jj}}{\partial Q}\right) \left(\frac{\partial m_{jj}}{\partial B}\right) - \left(\frac{\partial^2 m_{jj}}{\partial Q \partial B}\right) \right]$$

$$-E\left(\frac{\partial^2 \ln L}{\partial B^2}\right) = \sum_k \sum_j n_k m_{jk} \left\{ \frac{1}{m_{jj}} \left[\frac{1}{m_{jj}} \left(\frac{\partial m_{jj}}{\partial B}\right)^2 - \left(\frac{\partial^2 m_{jj}}{\partial B^2}\right) \right] + \left[\frac{k-j}{B^2} \right] \right\}$$

LOG-LINEAR MODEL FOR SMALL POPULATION GROUPS

$$\pi_{ij} = \binom{j}{i} \pi_{ii} B^{(j-1)} Q^{j(j-1)}, \quad i < j$$

$$\text{LOG } \pi_{ij} - \text{LOG } \pi_{ii} - \text{LOG} \binom{j}{i} = (j-1) \text{LOG } B + j(j-1) \text{LOG } Q, \quad i < j$$

$$\text{Let } a_{ij} = \text{LOG} \binom{j}{i}, \quad \beta = \text{LOG } B, \quad \gamma = \text{LOG } Q, \\ (B = e^\beta), \quad (Q = e^\gamma)$$

and substituting we have

$$\text{LOG } \pi_{ij} - \text{LOG } \pi_{ii} - a_{ij} = (j-1) \beta + j(j-1) \gamma$$

$$A \text{ LOG } \pi - \underline{a} = X \underline{\beta}$$

Log-Linear Model

WEIGHTED LEAST-SQUARES

HABER, LONGINI, COTSONIS (1988)

PARAMETER ESTIMATION COMPARING PROBABILITY AND

LOG-LINEAR MODELS - INFLUENZA A(H3N2)

1980-81 TECUMSEH DATA

I. Probability Model Using Maximum Likelihood

Number of Susceptibles/Household

Number	1		2		3		4	
	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
0	44	47.6	62	65.3	47	41.1	38	37.4
1	10	6.4	13	14.0	8	10.6	11	10.3
2	-	-	9	4.7	2	5.7	7	6.6
3	-	-	-	-	3	2.6	5	4.9
4	-	-	-	-	-	-	1	2.8
Total	54	54.0	84	84.0	60	60.0	62	62.0

$$\chi^2_8 = 11.6 \quad p = .17$$

$$\hat{SAR} = 20.1 \pm 3.5 \quad , \quad \hat{CPI} = .118 \pm .013$$

II. Log-Linear Model Using Weighted Least Squares

$$\chi^2_8 = 9.9 \quad p = .27$$

$$\hat{SAR} = 22.8 \pm 2.0 \quad , \quad \hat{CPI} = .110 \pm .007$$

OBSERVED AND EXPECTED DISTRIBUTION OF INFLUENZA B
1977 - TECUMSEH DATA
NUMBER OF SUSCEPTIBLES/HOUSEHOLD

NUMBER INFECTED	1		2		3		4		5	
	Obs.	Exp.	Obs.	Exp.	Obs.	Exp.	Obs.	Exp.	Obs.	Exp.
0	42	44.1	79	71.4	23	26.4	16	17.8	6	5.5
1	14	11.9	24	36.9	24	19.6	19	16.9	7	6.2
2	-	-	12	6.7	7	6.9	8	8.5	4	4.0
3	-	-	-	-	0	1.1	2	2.5	1	1.7
4	-	-	-	-	-	-	1	.4	0	.5
5	-	-	-	-	-	-	-	-	0	.1
TOTAL	56	56.0	115	115.0	54	54.0	46	46.1	18	18.0

$$\hat{Q} = .960$$

$$\hat{B} = .788 \quad \hat{CPI} = .212$$

$$\text{VAR}(\hat{Q}) = .0007$$

$$\text{VAR}(\hat{B}) = .0003$$

$$\text{Cov}(\hat{Q}, \hat{B}) = -.0002$$

$$\chi^2(13) = 15.1$$

$$P = .30$$

ACTUAL SAR = 4.0%

APPARENT SAR = 22.6%

OBSERVED AND EXPECTED DISTRIBUTION OF RESPIRATORY SYMPTOMS
DURING H3N2 EPIDEMIC PERIOD OF 1977-78 IN TECUMSEH

NUMBER ILLNESSES	1		2		3		4		5	
	OBS.	EXP.	OBS.	EXP.	OBS.	EXP.	OBS.	EXP.	OBS.	EXP.
0	38	35.4	48	39.5	13	14.9	6	9.9	4	3.3
1	19	21.6	29	40.0	19	18.9	13	14.9	3	4.8
2			25	22.5	20	17.8	17	16.4	9	6.3
3					10	10.4	23	15.9	8	7.6
4							7	9.8	5	7.9
5									6	5.2
TOTAL	57	57.0	102	102.0	62	62.0	66	66.0	35	35.1

$\hat{Q} = 0.835 \pm 0.024$, ACTUAL SAR = 16.5 ± 2.4 , APPARENT SAR = 46.7,

$\hat{B} = 0.622 \pm 0.020$; $\hat{CPI} = 0.378 \pm 0.020$, $Cov(\hat{Q}, \hat{B}) = -0.0002$,

$\chi^2(13) = 14.95$, P VALUE = 0.31, N = 322.

OBSERVED AND EXPECTED DISTRIBUTION OF INFLUENZA A(H1N1) 1978-1979
TECUMSEH DATA

NUMBER INFECTED	1		2		3	
	OBS.	EXP.	OBS.	EXP.	OBS.	EXP.
0	28	34.1	22	18.7	4	2.3
1	33	26.9	18	22.9	3	3.2
2			20	18.4	5	4.0
3					1	3.5
TOTAL	61	61.0	60	60.0	13	13.0

$\hat{SAR} = 22.5 \pm 8.6$, APPARENT SAR = 48.2, $\hat{CPI} = 0.441 \pm 0.038$.

$\chi^2(4) = 7.62$, P = .106

PARAMETERS

r - level of susceptibility ($r = 1, 2, \dots, R$)

h - level of infectiousness ($h = 1, 2, \dots, H$)

B_r = probability of escape from the community

$CPI_r = 1 - B_r$ community probability of infection

Q_{rh} = probability of escape from infected household member

$SAR_{rh} = (1 - Q_{rh}) \times 100$ secondary attack rate

DATA STRUCTURE

Household with s initial susceptibles

Susceptibility $r = (r_1, \dots, r_s)$

Infectiousness $h = (h_1, \dots, h_s)$

Infection $x = (x_1, \dots, x_s)$

where $x_j = \begin{cases} 0 & \text{if } j\text{-th person not infected,} \\ 1 & \text{if } j\text{-th person infected.} \end{cases}$

k_h - number persons infected at h -th level

$$k = \sum_h k_h$$

Find $P[x|r,h] : 2^s \times R^s \times H^s$

e.g., $R = 2, H = 2, s = 5$

$$2^5 \times 2^5 \times 2^5 = 32,768$$

PROBABILITY MODEL

$$p(\mathbf{x}|\mathbf{r},\mathbf{h}) = \begin{cases} \prod_{i=1}^s B_{r_i} & \text{for } k = 0 \\ p(1_k | \mathbf{r}_j, \mathbf{h}_j) \prod_{i: x_i=0} B_{r_i} \prod_{h=1}^H [Q_{r_i h}]^{k_h} & \text{for } k=1, \dots, s-1 \\ 1 - \sum_{\mathbf{x}: \sum x_i < k} p(\mathbf{x}|\mathbf{r},\mathbf{h}) & \text{for } k = s \end{cases}$$

where $\mathbf{j} = (j_1, \dots, j_k)$ are the indices of the k infected individuals.

$\mathbf{r}_j = (r_{j_1}, \dots, r_{j_k})$ and $\mathbf{h}_j = (h_{j_1}, \dots, h_{j_k})$. Also, 1_k denotes the array $(1, \dots, 1)$ of order k .

Constructed recursively -- e.g., $R = 2, H = 2, s = 5$

There are 37,448 probabilities.

EXAMPLE

$$R = 2, \quad S = 1, 2, \quad H = 1$$

WHEN $S = 1$: $2^1 \times 2^1 = 4$ PROBABILITIES

$$P(0|1) = B_1, \quad P(1|1) = 1 - B_1$$

$$P(0|2) = B_2, \quad P(1|2) = 1 - B_2$$

WHEN $S = 2$: $2^2 \times 2^2 = 16$ PROBABILITIES

$$P(0,0|1,1) = B_1^2, \quad P(0,0|1,2) = P(0,0|2,1) = B_1 B_2$$

$$P(0,0|2,2) = B_2^2$$

$$P(1,0|1,1) = B_1 Q_1, \quad P(1|1) = B_1 (1 - B_1) Q_1 = P(0,1|1,1)$$

$$P(1,0|1,2) = B_2 Q_2, \quad P(1|1) = B_2 (1 - B_1) Q_2 = P(0,1|2,1)$$

$$P(1,0|2,1) = B_1 Q_1, \quad P(1|2) = B_1 (1 - B_2) Q_1 = P(0,1|1,2)$$

$$P(1,0|2,2) = B_2 Q_2, \quad P(1|2) = B_2 (1 - B_2) Q_2 = P(0,1|2,2)$$

$$P(1,1|1,1) = 1 - B_1^2 - 2 B_1 (1 - B_1) Q_1$$

$$P(1,1|1,2) = 1 - B_1 B_2 - 2 B_2 (1 - B_1) Q_2 = P(1,1|2,1)$$

$$P(1,1|2,2) = 1 - B_2^2 - 2 B_2 (1 - B_2) Q_2$$

Infection attack rates by pre-season hemagglutination inhibition (HI) titer level stratified by age group: Influenza A(H3N2) epidemic seasons (1977-1978) and (1980-1981) combined in Tecumseh, Michigan

Pre-HI titer (1:x)	<u>Infection status</u>			Attack Rate [†]	Risk Ratio ^{††}
	No. infected	No. not infected	Total		
Children (0-17)					
Low level (x<8)	100	200	300	0.333	3.330*
Higher level (8≤x≤64)	20	180	200	0.100	
Total	120	380	500^{***}	0.240	
Adults (18+)					
Low level (x<8)	96	440	536	0.179	1.884*
Higher level (8≤x≤64)	42	402	444	0.095	
Total	138	842	980^{***}	0.141	

† Attack rate = No. infected/No. at risk

†† Risk ratio = Ratio of the attack rates

* p < 0.001

*** The total of 1480 individuals does not include the 26 "immune" individuals

The risk ratios across levels of age or across levels of pre-season HI tite are different (p < 0.0001) using the chi-square test for lack of interaction.

STATISTICS NOT APPROPRIATE

Observed and expected frequencies for households by the number of susceptibles from the influenza A(H3N2) seasons (1977-1978) and (1980-1981) combined,

in Tecumseh, Michigan, stratified by pre-season HI titer level[†]

No. susceptible/household Pre-season titer level		No. infected/household Pre-season titer level		No. of households		
low	higher	low	higher	observed	expected	
1	0	0	0	63 {	45	52.7
		1	0		18	10.3
0	1	0	0	70 {	65	63.6
		0	1		5	6.4
2	0	0	0	71 {	52	49.6
		1	0		11	14.4
		2	0		8	7.0
1	1	0	0	66 {	52	50.1
		0	1		2	3.8
		1	0		8	9.6
		1	1		4	2.5
0	2	0	0		45	42.9
		0	1		6	8.5(a)
		0	2		1	0.6(a)
3	0	0	0		17	16.9
		1	0		4	5.5
		2	0		3	3.9
		3	0		5	2.7
2	1	0	0		28	24.7
		0	1		1	1.3(b)
		1	0		6	7.0
		1	1		0	1.4(b)
		2	0		2	3.3
		2	1		2	1.2(b)
1	2	0	0		16	17.2
		0	1		6	2.5(c)
		0	2		0	0.1(c)
		1	0		2	3.2
		1	1		1	1.7(c)
		1	2		0	0.2(c)

0	3	0	0	11	11.2
		0	1	4	3.3(d)
		0	2	0	0.5(d)
		0	3	0	0.0(d)
4	0	0	0	16	13.7
		1	0	4	4.3
		2	0	6	3.5
		3	0	0	3.5
		4	0	2	3.0
3	1	0	0	13	13.8
		0	1	0	0.6(e)
		1	0	6	4.3
		1	1	1	0.6(e)
		2	0	1	3.0
		2	1	0	0.8(e)
		3	0	5	2.0(e)
		3	1	0	0.8(e)
2	2	0	0	11	11.5
		0	1	0	1.3(f)
		0	2	1	0.1(f)
		1	0	1	3.2
		1	1	3	1.2(f)
		1	2	1	0.1(f)
		2	0	3	1.5(f)
		2	1	0	1.0(f)
		2	2	0	0.2(f)
1	3	0	0	10	12.5
		0	1	5	2.7
		0	2	0	0.3(g)
		0	3	0	0.0(g)
		1	0	2	2.3(g)
		1	1	1	1.7(g)
		1	2	2	0.4(g)
		1	3	0	0.0(g)
0	4	0	0	10	8.2
		0	1	2	3.1(h)
		0	2	0	0.7(h)
		0	3	0	0.1(h)
		0	4	0	0.0(h)

5	0	0	0	3	2.4
		all other		3	3.6
4	1	0	0	2	2.7
		all other		4	3.3
3*	2*			4	-
2	3	0	0	4	5.2
		all other		6	4.8
1*	4*			2	-
0*	5*			3	-
Total				567	

Point estimates and standard errors:

Low pre-season antibody titer $\hat{CPI}_1 = 0.164 \pm 0.015$ $\hat{SAR}_1 = 26.0 \pm 3.0$

Higher pre-season antibody titer $\hat{CPI}_2 = 0.092 \pm 0.013$ $\hat{SAR}_2 = 2.1 \pm 2.6$

Overall goodness-of-fit $\chi^2(28df)^* = 33.01, p = 0.235$

Tests: $H_0 : CPI_1/CPI_2 = 1$ $\hat{RR} = 1.783$ $p < 0.0005$

$H_0 : SAR_1/SAR_2 = 1$ $\hat{RR} = 12.4$ $p < 0.0001$

† Only households with 5 or fewer susceptibles are included.

(a)-(h) Outcomes with the same letter were pooled for the goodness-of-fit test.

* These combinations were not included in the goodness-of-fit test.

INFLUENZA A(H3N2) CONTINUED

VARIABLE SUSCEPTIBILITY: PRE-SEASON ANTIBODY \times AGE

r = 1,2,3,4.

1. LOW, 0 - 17

2. HIGHER, 0 - 17

3. LOW, 18⁺

4. HIGHER, 18⁺

Estimated CPIs and SARs

Age	Pre-season titer				RR
		$X < 8$		$8 \leq X \leq 64$	Low to high titer
Children	CPI ₁	0.23 ± 0.03	CPI ₂	0.09 ± 0.03	2.5**
	SAR ₁	36.6 ± 6.2	SAR ₂	3.4 ± 4.7	10.8**
Adults	CPI ₃	0.13 ± 0.02	CPI ₄	0.09 ± 0.01	1.5
	SAR ₃	18.2 ± 4.4	SAR ₄	1.6 ± 3.7	11.4**

Goodness-of-fit $\chi^2 = 24.8$ (df = 23), $p = 0.36$, ** $p < 0.05$

ANTIBODY EFFICACY

DIRECT MEASURES

$$AE_{S,CI} = 1 - \frac{AR_H}{AR_L} \quad AR \quad \text{LEV. IV}$$

$$AE_{S,CI} = 1 - \frac{CPI_H}{CPI_L} \quad CPI \quad \text{LEV. IV}$$

$$AE_{S,P} = 1 - \frac{SAR_H}{SAR_L} \quad SAR \quad \text{LEV. I}$$

ESTIMATED ANIBODY EFFICACY

INDEX	CHILDREN	ADULTS	CRUDE
AR	0.700	0.469	0.588
CPE	0.593	0.321	0.439
SAR _{LK}	0.907	0.912	0.919

* **Table 3**
Maximum likelihood estimates and standard errors for parameters of the model of influenza A(H3N2) infections in 1977-1978 and 1980-1981 combined epidemics in Tecumseh, Michigan, with unrestricted contact parameters by age (0-17 vs 18+)

		Estimate	Transformation
Constant distribution: $T_1 \equiv 4.1$			
		$^a\beta_{11} = .0805 \pm .0208$	$SAR_{11} = 28.1186 \pm 6.1227$
		$\beta_{12} = .0354 \pm .0291$	$SAR_{12} = 13.4996 \pm 10.314$
		$\beta_{21} = .0268 \pm .0135$	$SAR_{21} = 10.4080 \pm 4.9593$
		$\beta_{22} = .0401 \pm .0127$	$SAR_{22} = 15.1662 \pm 4.4096$
Child		$B_1 = .8184 \pm .0254$	$CPI_1 = .1816 \pm .0254$
Adult		$B_2 = .8897 \pm .0128$	$CPI_2 = .1103 \pm .0128$
	Log likelihood	$= -522.333$	

* ADDY, ET AL. BIOMETRICS 47, 961-974 (1991)

The Transmissibility and Control of Pandemic Influenza A (H1N1) Virus

Yang Yang,¹ Jonathan D. Sugimoto,^{1,2} M. Elizabeth Halloran,^{1,3} Nicole E. Basta,^{1,2} Dennis L. Chao,¹ Laura Matrajt,⁴ Gail Potter,⁵ Eben Kenah,^{1,3,6} Ira M. Longini Jr.^{1,3*}

¹Center for Statistics and Quantitative Infectious Diseases, Fred Hutchinson Cancer Research Center and the University of Washington, Seattle, WA, USA. ²Department of Epidemiology, School of Public Health, University of Washington, Seattle, WA, USA. ³Department of Biostatistics, School of Public Health, University of Washington, Seattle, WA, USA. ⁴Department of Applied Mathematics, University of Washington, Seattle, WA, USA. ⁵Department of Statistics, University of Washington, Seattle, WA, USA. ⁶Department of Global Health, University of Washington, Seattle, WA, USA.

*To whom correspondence should be addressed. E-mail: longini@scharp.org

Pandemic influenza A (H1N1) 2009 (pandemic H1N1) is spreading throughout the planet. It has become the dominant strain in the southern hemisphere, where the influenza season is underway. Here, based on reported case clusters in the USA, we estimate the household secondary attack rate for pandemic H1N1 to be 27.3% [95% confidence interval (CI) 12.2%–50.5%]. From a school outbreak, we estimate a school child infects 2.4 (95% CI: 1.8–3.2) other children within the school. We estimate the basic reproductive number, R_0 , to range from 1.3 to 1.7 and the generation interval to range from 2.6 to 3.2 days. We use a simulation model to evaluate the effectiveness of vaccination strategies in the USA for Fall 2009. If vaccine were available soon enough, vaccination of children, followed by adults, reaching 70% overall coverage, in addition to high risk and essential workforce groups, could mitigate a severe epidemic.

corresponding to the rainy season. The last influenza pandemic was the Hong Kong A (H3N2) 1968–1969 pandemic. At that time, the first large epidemic was in Hong Kong in July 1968, followed by epidemics in South East Asia in August–September 1968, in the upper northern hemisphere between September 1968 and April 1969 (peaking in late December 1968 and early January 1969) and in the lower southern hemisphere between June and September 1969 (9). In the USA and the upper northern hemisphere, shifted (i.e., pandemic) or drifted strains of influenza tend to have a relatively small Spring “herald wave” before returning in the Fall (10). In the upper northern hemisphere, the 1918–1919 A (H1N1) pandemic had a mild Spring 1918 herald wave, followed by a severe second wave in the Fall of 1918. Pandemic Asian influenza A (H2N2), 1957–1958, caused mid-Summer 1957 outbreaks in Louisiana schools that were open in the Summer because of the need for children helping

Estimating Transmissibility

- Natural history of disease
 - Incubation period: $\Pr(\delta = l) = \alpha_l$, subject to $\sum_{l=\delta_{\min}}^{\delta_{\max}} \alpha_l = 1$
 - Infectious period: from ILI onset \tilde{t}_i to $\tilde{t}_i + D$, and probability of being infectious is β_d for day $\tilde{t}_i + d - 1$.
- Transmission model
 - Daily transmission probabilities: b (C2P) and $p(t)$ (P2P)
 - Daily escape probability

$$e_i(t) = (1 - b) \prod_{j \in h(i)} (1 - p(t) \beta_{t - \tilde{t}_j + 1})$$

$$L_i(b, p \mid \tilde{\mathbf{t}}_h) = \begin{cases} \prod_{t=1}^T e_i(t), & \tilde{t}_i = \infty, \\ \sum_{t=\tilde{t}_i - \delta_{\max}}^{\tilde{t}_i - \delta_{\min}} \alpha_{\tilde{t}_i - t} (1 - e_i(t)) \prod_{\tau=1}^t e_i(t), & \tilde{t}_i < T \end{cases}$$

Estimating Transmissibility (continue)

- Accounting for missing data
 - Household sizes and some ILI onset dates are missing
 - Multiple imputation (Schaffer, 1997).

- Calculating SAR and R_0 for US data

$$SAR = 1 - \prod_{d=1}^D (1 - p\beta_d)$$

$$R_0 = f(R_H + R_C + R_S) + (1 - f)(R_H + R_C)$$

$$R_H = N_H \times SAR_H, \quad R_S = N_S \times SAR_S, \quad R_C = \rho R_H$$

- Calculating R_0 for Mexican data
 - For large population, chain binomial becomes Poisson.
 - Assume each case corresponds to $K-1$ uninfected people.

Household SAR

	Individuals 11 and 12 are infected			Individuals 11 and 12 are not infected		
	Estimates	s.e.	95% CI	Estimates	s.e.	95% CI
Individual 10 is an index case						
p_H	0.0625	0.0268	(0.0265, 0.140)	0.0457	0.0213	(0.0181, 0.111)
SAR_H	0.273	0.101	(0.122, 0.505)	0.207	0.0870	(0.0849, 0.425)
R_H	0.82	0.303	(0.397, 1.693)	0.622	0.261	(0.274, 1.416)
Individual 10 is a secondary case						
p_H	0.0709	0.0288	(0.0314, 0.152)	0.0543	0.0232	(0.0231, 0.122)
SAR_H	0.304	0.105	(0.143, 0.535)	0.242	0.0915	(0.107, 0.459)
R_H	0.913	0.314	(0.466, 1.791)	0.726	0.274	(0.346, 1.523)

Household SAR: Sensitivity

		Incubation Period			Infectious Period		
		Estimates	s.e.	95% CI	Estimates	s.e.	95% CI
Short [†]	p_H	0.0447	0.0208	(0.0177, 0.108)	0.0662	0.0305	(0.0262, 0.157)
	SAR_H	0.204	0.0854	(0.0834, 0.418)	0.203	0.0853	(0.0833, 0.417)
	R_H	0.611	0.256	(0.268, 1.390)	0.610	0.256	(0.268, 1.388)
Long [†]	p_H	0.0462	0.0216	(0.0183, 0.112)	0.0376	0.0176	(0.0149, 0.0919)
	SAR_H	0.210	0.0880	(0.0857, 0.429)	0.211	0.0884	(0.0862, 0.431)
	R_H	0.629	0.264	(0.276, 1.432)	0.632	0.265	(0.278, 1.438)


[†] Short incubation period: $\delta_{min} = 1$, $\delta_{max} = 3$ and $(\alpha_1, \alpha_2, \alpha_3) = (0.6, 0.3, 0.1)$

Long incubation period: $\delta_{min} = 1$, $\delta_{max} = 4$ and $(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = (0.25, 0.25, 0.25, 0.25)$

Short infectious period: $D = 7$ and $(\beta_1, \dots, \beta_7) = (1.0, 1.0, 0.6, 0.4, 0.2, 0.1, 0.05)$

Long infectious period: $D = 8$ and $(\beta_1, \dots, \beta_8) = (1.0, 1.0, 1.0, 1.0, 1.0, 0.7, 0.4, 0.1)$

RESULTS: Historic Influenza SAR Estimates



RESULTS: Historic Influenza SAR Estimates

		Article	Cases SAR	Type of Confirmed	Data Source
A (H1N1)	2009	this paper	27.3 [12.2, 50.5]	ILL	U.S.
A (H5N1)	2006	(19) Yang 2007	29 [15, 51]	ILL-LAB	Sumatra
A (H2N2)	1957 (pandemic)	(20) Nishiura 2007	7.06 [4.73-9.44]	ILL	Tokyo
	1957 (pandemic)	(20) Nishiura 2007	9.07 [6.73-11.53]	ILL	Osaka
	1957 (pandemic)	(21) Longini 1982	16.6 [1, 32.5] (hh3)	ILL	Sugiyama
A (H1N1)	pre 1978	(21) Longini 1982	39.9 (hh4)	ILL	Hope, Simpson, Sutherland
	1978-79 ^a	(22) Longini 1982	30.6 [21.9, 39.3]	LAB	Seattle
	1978-79	(21) Longini 1982	33.6 (hh5)	ILL	Hope, Simpson, Sutherland
A (H3N2)	1977-78	(22) Longini 1982	14.7 [11, 18.4]	LAB	Tecumseh
	1978-79	(22) Longini 1982	20.6 [16.5, 24.7]	LAB	Seattle
	1977-78, 80-81	(23) Longini 1988	26 [23, 29]	LAB	Tecumseh
	1999-2000	(24) Viboud 2004	18.1	ILL	France
	1999-2000	(25) Cauchemez 2004	43 [39, 48] (hh2)	ILL	France
	1999-2000	(25) Cauchemez 2004	31 [28, 36] (hh3)	ILL	France
	1999-2000	(25) Cauchemez 2004	25 [22, 27] (hh4)	ILL	France
B	1961 ^b	(20) Nishiura 2007	18.41 [11.37, 25.95]	ILL	Osaka
	1961	(20) Nishiura 2007	10.51 [8.01, 13.15]	ILL	Osaka
	1975-76	(22) Longini 1982	12.7 [7.8, 17.6]	LAB	Seattle
Mixed	1998-1999 and 2000-2001 ^c	(28) Yang 2009 ^d	12.6 [6.5, 23.2]	ILL-LAB	
	1998-1999 and 2000-2001 ^e	(31) Halloran 2007	17.5 [12.6, 23.2]	ILL-LAB	
	1998-1999 and 2000-2001 ^f	(32) Yang 2007	12.8 [9.9, 15.7]	ILL-LAB	
	1998-1999 and 2000-2001 ^g	(11) Yang 2006	8.9 [6.4, 11.4]	ILL-LAB	
	2007 ^h	(33) Cowling 2008	6 [3, 10]	LAB	Hong Kong trial
	2007	(33) Cowling 2008	18 [13, 24]	ILL	Hong Kong trial
	2007	(33) Cowling 2008	11 [7, 16]	ILL	Hong Kong trial
2007	(33) Cowling 2008	4 [2, 6]	ILL	Hong Kong trial	