

Hypothesis Testing

Session 6

Module 1 Probability & Statistical Inference

The Summer Institutes

DEPARTMENT OF
BIostatISTICS

SCHOOL OF PUBLIC HEALTH

UNIVERSITY *of* WASHINGTON



Hypothesis Testing: Example questions

1. Is the chance of getting a cold different when participants take vitamin C than when they take placebo?
2. Suppose that 6 out of 15 grade-school students in a class develop influenza, whereas 20% of grade-school children nationwide develop influenza.

Is there evidence of an excessive number of cases in the class?

Session 6

PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY of WASHINGTON



Hypothesis Testing Example

In a study of 49 pregnant women living with HIV, mean arterial pressure is 79.4 mm Hg.

In the general population of pregnant women, the mean arterial pressure is 78.8 mm Hg with standard deviation of 1.4 mm Hg.

Are the data from our study consistent with the data from the general population?

Session 6

PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY *of* WASHINGTON



Selecting the Hypotheses to Test

Define:

μ = *population* mean (e.g. arterial pressure at 36 weeks gestation for pregnant women living with HIV)

Hypotheses:

Null Hypothesis: Generally, the hypothesis that the unknown parameter equals some fixed value.

$$H_0: \mu = 78.8 \text{ mm Hg}$$

Alternative Hypothesis: Contradicts the null hypothesis.

$$H_A: \mu \neq 78.8 \text{ mm Hg}$$

Session 6

PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY of WASHINGTON



Quantifying correctness

α = size or Type 1 error

$1 - \beta$ = power or $1 -$ Type 2 error

We assume that either H_0 or H_A is true.

Based on the data, we will choose one of these hypotheses.

		Truth	
		H_0 Correct	H_A Correct
Decision	Fail to reject H_0	$1 - \alpha$	β
	Reject H_0 , Accept H_A	α	$1 - \beta$


Setting the type 1 error

Set size/Type 1 error of the test, α . e.g., $\alpha = 0.05$.

$$\begin{aligned} 0.05 &= \alpha = P[\text{choose } H_A \mid H_0 \text{ true}] \\ &= \alpha = P[\text{reject } H_0 \mid H_0 \text{ true}] \end{aligned}$$

Q: How to construct a procedure that makes this error with only 0.05 probability?

A: Suppose we assume H_0 is true and suppose that, using that assumption, the data should give us a standard normal, Z .



If $\mu = 0$ then $|Z|$ is rarely large. A large $|Z|$ would make us question whether $\mu = 0$.

Rejecting the null hypothesis

Therefore, **we reject** H_0 if $|Z| > 1.96$.

$$\alpha = P[\text{reject } H_0 \mid H_0 \text{ true}] = 0.05$$

Then if we do find a large value of $|Z|$ we can claim that:

Either H_0 is true and something unusual happened (with probability α)...

or H_0 is not true.

But note that we operate under the assumption that H_0 is true and look for evidence suggesting it is false. We can reject the null but we can't prove it is true. Therefore, we say:

- If $|Z| > 1.96$, then we **reject** H_0
- If $|Z| < 1.96$, then we **fail to reject** H_0 .



Hypothesis Testing Example

Mean Arterial Pressure Example:

Let μ be the mean arterial pressure at 36 weeks for pregnant women living with HIV. In our sample of 49 women:

$$\bar{X} = 79.4 \text{ mm Hg}$$

For the general population of pregnant women at 36 weeks:

μ_0 = mean arterial pressure = 78.8 mm Hg

σ = std. dev. of mean arterial pressure = 1.4 mm Hg

Null hypothesis (H_0): mean for pregnant women living with HIV (μ) is the same as the mean for the general population of pregnant women (μ_0).

Alternative hypothesis (H_A): mean for pregnant women living with HIV (μ) is different than the mean for the general population of pregnant women (μ_0).

$$H_0 : \mu = \mu_0 = 78.8 \text{ mm Hg}$$

$$H_A : \mu \neq \mu_0 \text{ (} \mu \neq 78.8 \text{ mm Hg)}$$

Session 6

PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY OF WASHINGTON



Hypothesis Testing Example

Mean Arterial Pressure Example:

Test H_0 with significance level $\alpha = 0.05$.

Under H_0 :

$$\frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} \sim N(0, 1)$$

Therefore,

• **Reject H_0 if** $\left| \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} \right| > 1.96$ gives an $\alpha = 0.05$ test.

• **Reject H_0 if** $\bar{X} > \mu_0 + 1.96 \frac{\sigma}{\sqrt{n}}$ or

$$\bar{X} < \mu_0 - 1.96 \frac{\sigma}{\sqrt{n}}$$

Session 6

PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY of WASHINGTON



Hypothesis Testing Example

cont'd

Test: **Reject** H_0 if $\bar{X} > 78.8 + 1.96 \frac{1.4}{\sqrt{49}}$
or $\bar{X} < 78.8 - 1.96 \frac{1.4}{\sqrt{49}}$

or $\bar{X} > 79.192$
 $\bar{X} < 78.408$

$\bar{X} = 79.4$ therefore we reject the null

Consider Z:

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}}$$


and **reject** H_0 if $Z < -1.96$ or $Z > 1.96$

Session 6

PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY of WASHINGTON





Session 6
PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY of WASHINGTON



Hypothesis Testing and the p-value

p-value:

- a measure of how well the data agree with the null hypothesis
- probability of obtaining a result as extreme or more extreme than the actual sample when H_0 is true
- **not** the probability that the null (or alternative) is true

Session 6

PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY of WASHINGTON



Hypothesis Testing and the P-value Example


Mean arterial pressure example

$$\bar{X} = 79.4 \text{ mm Hg}, \quad n = 49, \quad \sigma = 1.4 \text{ mm Hg}$$

$$H_0 : \mu = 78.8 \text{ mm Hg}$$

$$H_A : \mu \neq 78.8 \text{ mm Hg}$$

p-value is given by: $2 * P[\bar{X} > 79.4] = 0.0027$



Session 6
PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY of WASHINGTON



(Somewhat Outdated) Guidelines for Judging the Significance of p-value

If $.05 \leq p < .10$, then the results are *marginally significant*.

If $.01 \leq p < .05$, then the results are *significant*.

If $.001 \leq p < .01$, then the results are *highly significant*.

If $p < .001$, then the results are *very highly significant*.

If $p > .1$, then the results are considered *not statistically significant* (sometimes denoted by NS).

Significance is not everything!

Session 6

PROBABILITY AND
CENTRAL STATISTICS

UNIVERSITY of WASHINGTON



American Statistical Association Guidelines for the Use of P-values

In March 2016, the ASA provided these guidelines:

1. P-values can indicate how incompatible the data are with a specified statistical model.
1. P-values do not measure the probability that the studied hypothesis is true, or the probability that the data were produced by random chance alone.
1. Scientific conclusions and business or policy decisions should not be based only on whether a p-value passes a specific threshold.
1. Proper inference requires full reporting and transparency.
1. A p-value, or statistical significance, does not measure the size of an effect or the importance of a result.
1. By itself, a p-value does not provide a good measure of evidence regarding a model or hypothesis.

Session 6

PROBABILITY AND
INFERENCE

UNIVERSITY of WASHINGTON



Hypothesis Testing and Confidence Intervals

Hypothesis Test: Fail to reject H_0 if:

$$\bar{X} < \mu_0 + Q_Z^{1-\frac{\alpha}{2}} \times \frac{\sigma}{\sqrt{n}}$$

and

$$\bar{X} > \mu_0 - Q_Z^{1-\frac{\alpha}{2}} \times \frac{\sigma}{\sqrt{n}}$$

Confidence Interval: Plausible values for μ are given by:

$$\mu < \bar{X} + Q_Z^{1-\frac{\alpha}{2}} \times \frac{\sigma}{\sqrt{n}}$$

and

$$\mu > \bar{X} - Q_Z^{1-\frac{\alpha}{2}} \times \frac{\sigma}{\sqrt{n}}$$

Session 6

PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY of WASHINGTON



1 vs 2-sided Hypothesis Testing

Depending on the alternative hypothesis, a test may have a **one-sided alternative** or a **two-sided alternative**. Consider

$$H_0 : \mu = \mu_0$$

We can envision (at least) three possible alternatives

$$H_A : \mu \neq \mu_0 \quad (1)$$

$$H_A : \mu < \mu_0 \quad (2)$$

$$H_A : \mu > \mu_0 \quad (3)$$

(1) is an example of a “two-sided alternative”

(2) and (3) are examples of “one-sided alternatives”

The distinction impacts:

- Rejection regions
- p-value calculation

Session 6

PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY of WASHINGTON



1-sided Hypothesis Testing

Mean Arterial Pressure Example: Instead of the two-sided alternative considered earlier we may have only been interested in the alternative that pregnant women living with HIV have higher mean arterial pressure.

$$H_0 : \mu = 78.8$$

$$H_A : \mu > 78.8$$

Given this, an $\alpha = 0.05$ test would reject when

$$\frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} = Z > Q_Z^{(1-0.05)} = 1.64$$

The p-value would be half of the previous,

$$\begin{aligned} \text{p-value} &= P[\bar{X} > 79.4] \\ &= 0.0014 \end{aligned}$$

Session 6
PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY of WASHINGTON



Hypothesis Testing Steps Summary

1. Identify H_0 and H_A
2. Identify a test statistic
3. Determine a significance level, $\alpha = 0.05$, $\alpha = 0.01$
4. Critical value determines rejection / acceptance region
5. Calculate p-value
6. Interpret the result

Session 6

PROBABILITY AND
INFERENCE STATISTICS

UNIVERSITY *of* WASHINGTON

