

Selection


Owing to this struggle for life, variations, however slight and from whatever cause proceeding, if they be in any degree profitable to the individuals of a species, in their infinitely complex relations to other organic beings and their physical conditions of life, will tend to the preservation of such individuals and will generally be inherited by the offspring. The offspring, also, will thus have a better chance of surviving, for, of the many individuals of a species which are periodically born, but a small number can survive. I have called this principle, by which each slight variation, if useful, is preserved, by the term Natural Selection.

Natural Selection


- More offspring produced that can survive/reproduce
- Individuals differ in ability to survive/reproduce (due to genotype)
- Genotypes that promote survival
 - Present in excess at reproductive age
 - Contribute disproportionately to offspring

Key points

- Selection acts on **individuals**
- Adaptation is a **population** process



Viability Selection



Model Assumptions

- Diploid individuals
- One locus (2 alleles)
- Sexual reproduction
- Discrete generations
- Random mating
- Infinite population size
- No mutation
- No migration
- Fitnesses constant

Fitness

- Individual: proportional contribution of offspring to next generation
- Genotype: average fitness for all individuals with that genotype in the population
- With VS, fitness is probability of survival to reproductive age

Absolute Fitness

2 alleles: A_1 (p), A_2 (q)

Genotypes	A_1A_1	A_1A_2	A_2A_2
Zygotic Frequencies	p^2	$2pq$	q^2
Absolute fitness	w_{11}	w_{12}	w_{22}
Adult Frequencies	p^2w_{11}	$2pqw_{12}$	q^2w_{22}

$$\bar{w} = p^2w_{11} + 2pqw_{12} + q^2w_{22}$$

Absolute Fitness

2 alleles: A_1 (p), A_2 (q)

Genotypes	A_1A_1	A_1A_2	A_2A_2
Zygotic Frequencies	p^2	$2pq$	q^2
Absolute fitness	w_{11}	w_{12}	w_{22}
Adult Frequencies	p^2w_{11}	$2pqw_{12}$	q^2w_{22}
Normalized Adult Frequencies	p^2w_{11}/\bar{w}	$2pqw_{12}/\bar{w}$	q^2w_{22}/\bar{w}

$$p' = \frac{p^2w_{11} + pqw_{12}}{\bar{w}}$$

$$\Delta_s p = p' - p = \frac{p^2w_{11} + pqw_{12} - p\bar{w}}{\bar{w}}$$

Relative Fitness

2 alleles: A_1 (p), A_2 (q)

Genotypes	A_1A_1	A_1A_2	A_2A_2
Zygotic Frequencies	p^2	$2pq$	q^2
Absolute fitness	w_{11}	w_{12}	w_{22}

Relative Fitness

2 alleles: A_1 (p), A_2 (q)

Genotypes	A_1A_1	A_1A_2	A_2A_2
Zygotic Frequencies	p^2	$2pq$	q^2
Absolute fitness	w_{11}	w_{12}	w_{22}
Relative fitness	1	w_{12}/w_{11}	w_{22}/w_{11}

Relative Fitness

2 alleles: A_1 (p), A_2 (q)

Genotypes	A_1A_1	A_1A_2	A_2A_2
Zygotic Frequencies	p^2	$2pq$	q^2
Absolute fitness	w_{11}	w_{12}	w_{22}
Relative fitness	1	w_{12}/w_{11}	w_{22}/w_{11}
Relative fitness	1	$1-hs$	$1-s$

s : Coefficient of selection

h : Dominance coefficient

$h = 0$: A_1 dominant

$h = 1$: A_1 recessive

$0 < h < 1$: incomplete dominance

$h = 0.5$: codominant

$h < 0$: overdominance

$h > 1$: underdominance

Relative Fitness

2 alleles: A_1 (p), A_2 (q)

Genotypes	A_1A_1	A_1A_2	A_2A_2
Zygotic Frequencies	p^2	$2pq$	q^2
Absolute fitness	w_{11}	w_{12}	w_{22}
Relative fitness	1	w_{12}/w_{11}	w_{22}/w_{11}
Relative fitness	1	$1-hs$	$1-s$
Adult frequencies	p^2	$2pq(1-hs)$	$q^2(1-s)$

$$\bar{w} = p^2 + 2pq(1-hs) + q^2(1-s)$$

Relative Fitness

2 alleles: A_1 (p), A_2 (q)

Genotypes	A_1A_1	A_1A_2	A_2A_2
Zygotic Frequencies	p^2	$2pq$	q^2
Absolute fitness	w_{11}	w_{12}	w_{22}
Relative fitness	1	w_{12}/w_{11}	w_{22}/w_{11}
Relative fitness	1	$1-hs$	$1-s$
Adult frequencies	p^2	$2pq(1-hs)$	$q^2(1-s)$
Normalized adult frequencies	p^2/\bar{w}	$2pq(1-hs)/\bar{w}$	$q^2(1-s)/\bar{w}$

Relative Fitness

2 alleles: A_1 (p), A_2 (q)

Genotypes	A_1A_1	A_1A_2	A_2A_2
Zygotic Frequencies	p^2	$2pq$	q^2
Absolute fitness	w_{11}	w_{12}	w_{22}
Relative fitness	1	w_{12}/w_{11}	w_{22}/w_{11}
Relative fitness	1	$1-hs$	$1-s$
Adult frequencies	p^2	$2pq(1-hs)$	$q^2(1-s)$
Normalized adult frequencies	p^2/\bar{w}	$2pq(1-hs)/\bar{w}$	$q^2(1-s)/\bar{w}$

$$p' = \frac{p^2 + pq(1-hs)}{\bar{w}}$$

$$\Delta_s p = p' - p = \frac{p^2 + pq(1-hs) - p\bar{w}}{\bar{w}}$$

Change in Allele Frequency



$$\Delta_s p = \frac{pq(s(ph + q(1-h)))}{\bar{w}}$$

$$(ph + q(1-h))$$



$$\Delta_s p > 0 \text{ if } s > 0$$

$$\Delta_s p < 0 \text{ if } s < 0$$


Dependency on s



Dependency on p



Dependency on h



Overdominance

2 alleles: A_1 (p), A_2 (q)

$$h < 0$$

$$W_{11} < W_{12}$$

$$W_{22} < W_{12}$$

Genotypes	A_1A_1	A_1A_2	A_2A_2
Zygotic Frequencies	p^2	$2pq$	q^2
Absolute fitness	W_{11}	W_{12}	W_{22}
Relative fitness	1	$1-hs$	$1-s$

Overdominance

2 alleles: A_1 (p), A_2 (q)


$$h < 0$$

$$W_{11} < W_{12}$$

$$W_{22} < W_{12}$$

Genotypes	A_1A_1	A_1A_2	A_2A_2
Zygotic Frequencies	p^2	$2pq$	q^2
Absolute fitness	W_{11}	W_{12}	W_{22}
Relative fitness	1	$1-hs$	$1-s$
Relative fitness	$1-s$	1	$1-t$

Overdominance



Underdominance

2 alleles: A_1 (p), A_2 (q)

$$h > 1$$

$$W_{12} < W_{11}$$

$$W_{12} < W_{22}$$

Genotypes	A_1A_1	A_1A_2	A_2A_2
Zygotic Frequencies	p^2	$2pq$	q^2
Absolute fitness	W_{11}	W_{12}	W_{22}
Relative fitness	1	$1-hs$	$1-s$

Underdominance










Table 3. Summary of behavior of the one-locus two-allele viability model.

	<i>AA</i>	<i>Aa</i>	<i>aa</i>	
Directional selection				
Case 1	$w_{11} > w_{12} > w_{22}$			<i>A</i> becomes fixed
				
Case 2	$w_{11} < w_{12} < w_{22}$			<i>a</i> becomes fixed
				
Overdominance				
Case 3	$w_{11} < w_{12} > w_{22}$			Stable polymorphism
				
Underdominance				
Case 4	$w_{11} > w_{12} < w_{22}$			Unstable equilibrium
	